

Required Withholding Period for Vine Leaves Following Spraying with Pesticide

Britt Marianna Maestroni¹, Iyad Ghanem², Raymond Correll³, Amer Abu Alnaser², Marivil Islam¹, Veronica Cesio⁴, Horacio Heinzen⁴, Andrew Cannavan¹

¹Food and Environmental Protection Laboratory, International Atomic Energy Agency, Vienna, Austria

²Department of Molecular Biology and Biotechnology, Syrian Arab Republic Atomic Energy Commission, Damascus, Syria

³Biometry Hub, University of Adelaide, Adelaide, Australia

⁴Grupo Analisis Contaminantes Trazas, University of the Republic, Montevideo, Uruguay

Email address:

b.m.maestroni@iaea.org (B. M. Maestroni), eghanem@aec.org.sy (I. Ghanem), rho.environmentics@bigpond.com (R. Correll), abualnaser@gmail.com (A. A. Alnaser), m.islam@iaea.org (M. Islam), veronicacesio@gmail.com (V. Cesio), heihoracio@gmail.com (H. Heinzen), a.cannavan@iaea.org (A. Cannavan)

To cite this article:

Britt Marianna Maestroni, Iyad Ghanem, Raymond Correll, Amer Abu Alnaser, Marivil Islam, Veronica Cesio, Horacio Heinzen, Andrew Cannavan. Required Withholding Period for Vine Leaves Following Spraying with Pesticide. *Journal of Health and Environmental Research*. Vol. 4, No. 4, 2018, pp. 140-152. doi: 10.11648/j.jher.20180404.14

Received: October 25, 2018; **Accepted:** November 10, 2018; **Published:** December 17, 2018

Abstract: Vine leaves are consumed in many countries but little attention is paid to the residues left on them after the application of pesticides that help prevent pests and protect the grapes, the economically important target. Therefore, it is of utmost importance to study the dissipation of the pesticides applied to this crop to protect the consumers that also eat vine leaves. Dissipation kinetics of chlorpyrifos, chlorpyrifos-methyl, diazinon and dimethoate residues were studied in vine leaves grown under sunny conditions in Syria, using an ethyl acetate based sample preparation followed by GC-MS/MS determination. The dissipation rate for all doses applied followed first-order kinetics, with half-lives in grape leaves in the range of 2.9 – 3.9 days. At the recommended application dose, a withholding period of 8.9-37.1 days before consumption should be applied to meet current MRLs and minimise risks to consumers. The effectiveness in the reduction of pesticide loads in vine leaves through washing with either cold or hot water was dependant on the physicochemical properties of the studied pesticides. Hot water washing was very effective for dimethoate, a polar and water-soluble pesticide, with an effective reduction of 92% of the residue level; but no significant effect was observed for chlorpyrifos, the most apolar compound in this study.

Keywords: Vine Leaves, Dissipation Kinetics, Half-Life, Withholding Period

1. Introduction

Grape (*Vitis vinifera*) was estimated by Food and Agriculture Organisation of the United Nations to be the most widely cultivated fruit crop in the world [1]. The Mediterranean countries constitute the main area of cultivation of grapevines. While the main product are grapes, with most of the production destined for wine making, vine leaves are also important for cultural and nutritional reasons. Vine leaves have been used as a nutritious food in Greece and the Middle East for centuries and their popularity as a healthy food is increasing globally [2]. In countries such as

Syria, where vine leaves are widely consumed, it is important to assess human dietary exposure to residues of pesticides applied to vine leaves. As vine leaves may represent an important contribution to the Mediterranean diet, data on the fate of pesticides in vine leaves after application are essential for both the calculation of the theoretical maximum daily intake and the establishment of the Maximum Residues Limits (MRLs) [3]. Most studies on pesticide residues deal with the analysis of grapes or the transformation from vine to wine, as discussed in the review by Grimalt et al. [4]. To date only a few studies have been dedicated to residues in vine leaves and foliage in general [5-10]. Insecticides and

fungicides are applied to be effective against a wide range of insect pests, such as grape moth (*Lobesia botrana*), and fungal diseases such as downy mildew (*Plasmopora viticola*), powdery mildew (*Uncinula necator*) and gray mold (*Botrytis cinerea*), but their use has to be strictly regulated. The legal parameter with respect to pesticide residues, which determines whether a food product may be placed on the market, is the MRL expressed in mg/kg of product. This value is calculated from toxicological data such as Acceptable Daily Intake (ADI in mg/kg/day) and Food Daily Intake (FDI in kg/day), and agronomic data such as active dose and dissipation curves in the field. FAO and the World Health Organization (WHO), through their Joint Meeting on Pesticide Residues (JMPR) meetings evaluate pesticides and their residues in food to set MRLs for food commodities [11-12]. The occurrence of pesticides in the environment, as well as in vine leaves, remains an important challenge to address in Syria and elsewhere. The agricultural practices, including pesticide management practices, must be continually improved to meet it. It is important to generate analytical data on the actual levels of crop protection chemicals on crops and in the environment, and to consider these data with respect to the agricultural practices applied under local conditions to be able to optimize pesticide management. In assessing the impact of dietary exposure to pesticides, a number of parameters have to be taken into account and carefully evaluated, including MRLs, withholding times and dissipation rates and post-harvest management, amongst others. These parameters are obtained either experimentally or through modelling and they vary according to the pesticide, the type of crop and the prevailing environmental conditions. A combination of effects results in a reduction, over time, from the initial amount of active ingredient on crops after application. The dissipation rate (K_{diss}) is an important kinetic parameter used to calculate residual concentrations of pesticides in crops harvested for human or animal consumption [13]. The term dissipation is defined as an integrated process where several effects play a role, including volatilization, photodegradation, wash-off, leaching, hydrolysis, chemical and biological degradation, among others [13]. Dissipation rates are usually expressed as the pesticide half-life; the half-life being the time required for the pesticide residue level to fall to half of the initial concentration directly after application. Because of the large combinations of pesticides and crops, it is very important to estimate dissipation half-lives to contribute to risk assessments, for example when establishing MRLs for pesticides in various commodities. The dissipation rate for pesticides applied on a specific crop depends on several factors: the chemical formulation and application method, climatic conditions - especially rainfall and temperature, vapour pressure of the pesticides, and the potential for photodegradation and other chemical degradation [14]. This means that dissipation curves are very specific to local conditions in each growing area [15-20], and therefore generation of precise knowledge of pesticide degradation kinetics is very important to produce reliable data for

international bodies, such as Codex Alimentarius Commission, that set MRLs [21]. Because of the agronomic behaviour and differences among regions in the world, the same active ingredients may have different MRLs depending on the country and the climatic conditions [22]. Fantke et al. [13] systematically reviewed 811 scientific literature sources, and analysed 4513 dissipation half-lives of 346 pesticides measured in 183 plants. The authors emphasized that further experiments are needed to analyse pesticide-plant species combinations that so far have not been covered and to allow the use of prediction models. The fundamental reason is because modelling dissipation in plants is highly uncertain and the estimation of dissipation half-lives strongly relies on experimental field data. A study was initiated in Syria on the behaviour of specific organophosphate pesticides (OP) in the field under the current agricultural practices for grape vines. The compounds included in the study were chlorpyrifos, chlorpyrifos methyl, diazinon and dimethoate. OPs are extensively used due to their high insecticidal activity and relatively low persistence [5]. The dissipation rate of OPs is very fast [23-25]. The objective of the study was to establish the withholding period (WHP) for these specific OP pesticides used in vineyards in Syria. The WHP is the minimum period of time that must be allowed after pesticide application before the treated area or crop can be grazed, cut for fodder or harvested. Withholding periods vary for different pesticide/crop combinations. They help to ensure that residues in the treated crop will not exceed the maximum residue limits when the crop is placed on the market [26]. When being prepared as a food in the home (or by industry), vine leaves are first washed. The fate of pesticide residues on raw agricultural crops has been well studied and recently well reviewed by Amvrazi [27], Kaushic [28], Holland [29]. According to these studies the rinsability of a pesticide is not always correlated with its water solubility and different pesticides may be rinsed off commodities using different washing procedures and washing agents [27]. In the present study, after sampling, vine leaves were analysed unwashed, washed with cold water, and washed with boiling tap water to assess whether domestic washing can offer a practical method for decreasing the intake of pesticides for consumers.

2. Materials and Methods

2.1. Field Trials

2.1.1. Site Descriptions

The experimental trials were carried out during the summer of 2016 in two different vineyards located outside the city of Damascus, Syria. A local cultivar of table grape was used for testing. The Al Soujah (Yafour) area, located 23 km west of Damascus, is an arid zone where the vineyard is surrounded by olive trees and is an experimental station belonging to the Atomic Energy Commission of Syria (AECS). The Al-Hamah area is a private vineyard in a greener zone located 14 km north west of Damascus and close to a river (see Figures A1 and A2 in the Appendix).

Meteorological data were collected by an agrometeorological station in Damascus. During the field trials the average daily temperature was 26°C, the maximum temperature was 36°C, the average humidity was 27%, the average solar radiation was 209 w/m², the average wind speed was 11.1 km/h and there was no rain. Each growing area was divided into two zones. One zone was treated and the other one was left untreated to be used as a control. The vines did not receive any irrigation, or pesticide or fertilizer treatment throughout the growing season. Four different formulations of organophosphate pesticides (OPs) were purchased from a local supplier and applied at the concentrations indicated on the labels, using a 20 L hand-pump pesticide sprayer (in Al Soujah area) and an electronic sprayer (in Al-Hamah area). Care was taken to ensure that the vines were well covered

with the spraying mixture. The formulations applied to the vine areas, their label concentrations and the target application rate for each pesticide are presented in Table 1. Formulated forms of the pesticides dimethoate (400 mg/mL) and chlorpyrifos ethyl (480 mg/mL) were from AgriPest, diazinon 600 mg/mL was from Orient for Veterinary Pesticides and chlorpyrifos methyl (20%, 200 mg/mL) was from OvaGreen. At the Al Soujah site, 50 m² of vines trees were treated with 7 L of the pesticide formulation mixture, at the Al-Hamah site 100 m² of vines were sprayed with 100 L of the formulation mixture. In both cases the total volume of prepared pesticides was sprayed on the vines to deliver the amount of kg/Ha as shown in Table 1. Only one pesticide application was carried out at both locations at the end of the flowering stage.

Table 1. OP insecticides, their concentration and the amount sprayed on vine leaves in Al-Hamah and Al Soujah area.

Insecticide	Label concentration (mg/mL)	Amount of OP used for the mixture in Al Hamah (mL)	Amount sprayed in Al Hamah (kg/Ha)	Amount of OP used for the mixture in Al Soujah (mL)	Amount sprayed in Al Soujah (kg/Ha)
Dimethoate	400	50	2	7	0.56
Diazinon	600	50	3	7	0.84
Chlorpyrifos-methyl	200	200	4	280	11.2
Chlorpyrifos-ethyl	480	50	2.4	7	0.672

2.1.2. Sampling

For studying the dissipation of the four evaluated OP pesticides, vine leaves were collected at time 0 (one hour after spraying, when the spraying mixture had dried) and then after 1, 7, 14 and 21 days. Sufficient leaves were randomly collected from at least 5 vine trees to provide an overall sample weight of 500 g of vine leaves. The samples were labelled and immediately dispatched to the laboratory, where they were frozen and kept at -18°C until analysis.

2.2. Chemicals

All chemical analyses were performed at the Joint FAO/IAEA laboratories in Seibersdorf, Austria. All water used in this study was obtained from an in-house purification system (Milli Q, Millipore, USA). Certified pesticide analytical standards and triphenylphosphate (TPP, certified analytical standard) were purchased from Sigma Aldrich, Austria. Stock solutions of individual standards were prepared at 10 mg/mL and used in the preparation of a mixed stock solution at 25 ng/μL. The individual stock solution and the stock mixture were stored in amber screw-capped vials with septa in the dark at -20°C. Working standard solutions were prepared from the mixed stock solution according to requirements. Residue grade glacial acetic acid and ethyl acetate were purchased from Merck (Austria) and ascorbic acid was purchased from Sigma Aldrich (Austria). Acidified ethyl acetate was prepared by dissolving ascorbic acid (0.5 g) in milliQ water (10 mL), to which glacial acetic acid (10 mL) was added and the volume made up to 1L with ethyl acetate. The solution was stored in a screw-capped bottle with septum, wrapped with aluminium foil and kept in the dark at 4°C. Residue grade sodium hydrogen carbonate (NaHCO₃), anhydrous sodium sulphate (NaSO₄) and anhydrous

magnesium sulphate (MgSO₄) were purchased from Sigma Aldrich (Austria). Residue grade primary-secondary amine (PSA) sorbent was purchased from Varian (USA).

2.3. Sample Preparation

Upon arrival at the laboratory each sample was divided into three parts. One part was directly stored at -20°C pending analysis. The other two parts were washed by immersion for 5 seconds into either boiling water or tap water, air dried in the dark and stored -20°C pending analysis. Before analysis, the vine leaves were prepared by removing the stems and crushing the leaves to small particles using liquid nitrogen in a mortar and pestle. To ensure effective homogenization, the samples were further homogenized, while still frozen, using a common food chopper (Moulinex, 1000 W) for 1 minute.

2.3.1. Analytical Method

A validated method for the determination of chlorpyrifos, chlorpyrifos-methyl, diazinon, and dimethoate was applied for the analysis of the samples [30]. Analytical portions of 2 g of homogenised vine leaves were weighed into 50 mL labelled PTFE centrifuge tubes. Milli Q water (4 mL) was added and the samples left to soak for 30 min. Acidified ethyl acetate (4 mL) and the surrogate standard, TPP (200 μL of a 2 ng/μL solution) were added to each sample. The tubes were shaken vigorously by hand for 1 minute and then for 30 min on a horizontal shaker. After this step the first salt mixture (0.3334 g NaHCO₃ + 2 g NaSO₄) was added, the tubes were hand-shaken vigorously and the contents were thoroughly homogenized using an ultra-turrax homogenizer (IKA-T25, IKA, Germany) at maximum speed for 1 min. The homogenates were centrifuged for 5 min at approximately 12,600 g at 20°C. Two mL of the organic supernatant were

cleaned up in an Eppendorf tube of 15 mL by adding a second salt mixture (50 mg PSA + 300 mg MgSO₄), and vortexed for 30 seconds. They were then more intensively agitated on an orbital vortex mixer (Scientific industries) at maximum speed for 5 min. The tubes were then centrifuged for 5 min at approximately 12,600 g at 20°C. The extracts were filtered using a 0.2 µm syringe filter yielding a 0.5 g/mL sample extract. Finally, 100 µL of internal standard (sulfotep, 1 ng/µL) was added to 900 µL of each extract, including calibration standards, to reach a final volume of 1 mL and injected into the GC-MS/MS triple quadrupole system.

2.3.2. Apparatus and Analytical Conditions

The gas chromatography system used to analyse the pesticide residues consisted of an Agilent 7693 autosampler, 7890 B GC and 7000C GC-MS/MS triple quadrupole system (Agilent Technologies, Santa Clara, USA). Data acquisition and processing were implemented using Agilent Mass Hunter Quantitative Analysis B07.00 software. Two 15.0 m x 0.25 mm ID x 0.25 µm HP-5 ms ultra-inert columns (Agilent J&W, USA) were connected by an electronic pressure controller to enable a 5-minute post-run backflush. The samples were injected using a multimode injector inlet in splitless mode through an ultra-inert inlet liner with a glass wool plug (Agilent Technologies, Santa Clara, USA). The injection volume was 3 µL. The temperature programmable injector port was set up to an initial injection temperature of 70°C for 0.02 minutes and then ramped up to 320°C (at 850°C/min) for 5 minutes, then cooled down to 280°C at 20°C/min. The oven temperature was held at 70°C for 1 minute, then ramped to 150°C (at a rate of 50°C/min), increased to 200°C at 6°C/min followed by a final ramp to 280°C at 16°C/min. The total run time was 24 min. Helium

(99.999% purity) was used as the carrier gas and nitrogen (99.999% purity) as the collision gas. The transfer line was maintained at 280°C. The retention time lock setting (RTL) used chlorpyrifos-methyl as the locking compound at retention time of 10.83 min. The instrument worked at a constant flow (1.25 mL/min in column 1 and 1.45 mL/min in column 2). The ion source and the quadrupole analyser temperatures were fixed at 300°C/min and 150°C/min, respectively. The transitions and the collision energies used for detection are shown in the Appendix (*Table A1 of the Appendix*). A twenty-five time-segment method was created to obtain adequate sensitivity, and in each time segment dwell times were optimised to collect at least 12 points across a peak (cycles between 3 and 4). The solvent delay was 4 minutes.

2.3.3. Residues Analytical Determination

The analytical calibration was carried out using matrix matched standards calibration curves, covering the range 0.005-0.240 µg/g (2.5-120 µg/L) and using the bracketing calibration modality. When needed, the samples and the corresponding blanks, used for the preparation of the matrix matched calibration curves, were diluted with acidified ethyl acetate to meet the analytical range of 0.005-0.240 µg/g. Where more than a single estimate of the residue level was available (based on different dilutions required for assessment) an average estimate was used for that sample. The limit of quantitation (LOQ) and the precision data obtained during method validation are presented in the Appendix (*Table A2 of the Appendix*). Current EU and Codex MRLs for chlorpyrifos, chlorpyrifos-methyl, diazinon and dimethoate are given in Table 2. Values below the minimum reporting level or lowest calibrated level (LCL) were not included in the data set.

Table 2. European Union MRL for grape leaves and current Codex MRL for commodities similar to vine leaves, and selected properties of the three OP pesticides (from Lewis et al. [31]).

Pesticide	MRL-EU (mg/kg)	MRL-Codex Alimentarius (mg/kg) for similar commodities	Koc	Kow	Half-life on plants (days)	Photolysis half-life (days)
Chlorpyrifos	0.05	0.5 (grapes)	8151	50100	3.3	29.6
Chlorpyrifos-methyl	0.05	0.5 (grapes)	4645	10000	2.4	1.74
Diazinon	0.01	0.5 (lettuce leaf)	609	4900	2.4	50
Dimethoate	0.01	0.5 (peppers)	8	5.06	3.7	175

2.4. Statistical Methods

In most of the cases pesticide residue dissipation curves can be described mathematically by a first order decay model [32- 38]. Occasionally other mathematical models have to be applied to describe the decay [12]. Results obtained from the analysis of sampled vine leaves were plotted and statistically analysed using the statistical software R (R Core Team, 2017) [39]. In this study, where possible a first order decay model was used to describe the data according to Equation 1:

$$C_t = C_0 \times \exp(-k_{\text{diss}} \times t) \quad (1)$$

Where C_t is the residual pesticide concentration in the vine leaves (mg/kg) at time t , C_0 is the initial pesticide

concentration, in mg/kg on the harvested plant material immediately after pesticide application, t the time since spraying and k_{diss} is the dissipation constant, which is related to the rate of pesticide breakdown. Equation 1 is non-linear and implicitly assumes a constant variance along time. A logarithmic transformation is therefore applied resulting in Equation 2, which is linear:

$$\ln(C_t) = \ln(C_0) - k_{\text{diss}} \times t \quad (2)$$

Equation 2, was applied to fit the residue data obtained for all the pesticides. An alternative approach was required for diazinon. In the case of diazinon, a biphasic model was required to obtain homoscedasticity, i.e. homogeneity of variances. This was achieved by minimising the sum of

squares of the residuals on the logarithmically transformed data. Rearranging Equation 2 and solving for k_{diss} yields Equation 3:

$$k_{\text{diss}} = (\ln(C_0) - \ln(C_t)) / (t) \quad (3)$$

k_{diss} represents the slope of the regression line. From this, one can finally obtain the half-life ($t_{1/2}$), defined as the time needed to reduce the initial pesticide residue level to half of its initial value [33], and calculated from Equation 4:

$$t_{1/2} = \ln(2) / k_{\text{diss}} \quad (4)$$

For each pesticide there were two data series, based on the different study areas. Where these two series had different starting residue levels at the time of the first sampling, a convenient method for estimating the initial application residue level was to use the intercepts of the regressions models for each data series separately. The maximum intercept value was used further to estimate the half-life and the required WHP for that pesticide. The WHP was estimated as the time needed for the residues to dissipate to values equivalent to the MRL after pesticide application (time 0) using the established regression model. The Equation used was:

$$\text{WHP} = [(\text{intercept} - \ln(\text{MRL value})) / k_{\text{diss}}] \quad (5)$$

Paramasivam *et al.* [40] used Equation 5 to calculate the pre-harvest interval (PHI) defined as the maximum time (days) required for the residues to fall below the MRL. Confidence limits (CL) for the regression curves were obtained from Equation 6:

$$\text{CL} = X \pm t \text{ SE} \quad (6)$$

Where t is the Student's t value and SE is the standard error. The statistical model used to describe the data included

Table 3. Comparative summary of the withholding period (WHP) for chlorpyrifos, diazinon and dimethoate according to the Codex and EU MRL. Estimated confidence intervals (in days) are for unwashed leaves. Lower confidence level (LCL) and upper confidence level (UCL).

Pesticide	Estimate WHP (Codex)	LCL-UCL	Estimate WHP (EU)	LCL-UCL
Chlorpyrifos	26.0	22.2-29.5;	37.0	31.6-42.1
Diazinon	8.9	6.0-10.6	37.1	n/a
Dimethoate	19	18-20	33	31-34

3.1. Chlorpyrifos Methyl

No detectable residues were found for chlorpyrifos methyl (CPM). None of the samples analysed showed CPM residues above the limit of detection. This behaviour can be explained by photolysis, as shown by the photolysis half-life data in Table 2, and a fast dissipation rate of CPM under the local conditions in Syria. This is in line with previous findings about this pesticide for which the half-life ranged from 0.97 to 3.27 days in Italy [17]) or 0.9-2.6 days on different foliage [7]. The concentration of chlorpyrifos methyl applied at the Al-Hamah site was very small (200 mL of 20% pesticide

formulation dissolved in 50 L of water) compared to that used at the Al Soujah site (280 mL of 20% pesticide dissolved in 7 L of water). However, there is no information about the authenticity of the pesticide formulation. No statistical analysis could be carried out on the data relating to CPM.

additional terms to take into account the effect of the study areas, the effect of washing and their interactions. Where the results indicated that these terms were not significant, then the model was simplified and those additional terms deleted. Confidence limits for the WHP were obtained from the confidence limits of the regression slope, using a stepwise approach, at the point of intersection between the regression line and the MRL. The first step in estimating the confidence limits for the WHP was to estimate the standard error of the fitted line at the intersection point of the regression line with the MRL value (as shown in Figure 1). Finally, the standard error value (SE) was divided by the slope (with its sign) of the regression line to obtain the SE of the WHP. The confidence interval for the WHP was then obtained from Equation 6. The appraisal of the withholding period required to achieve the MRL value for diazinon was estimated using a two-compartment model following a Newton-Raphson technique [41]. Confidence limits of the required withholding period were estimated for the two-compartment model using a bootstrap technique [42], which was used to obtain 1000 estimates. The quantiles of the estimates obtained were used for the estimation of the confidence limits of the withholding period.

3. Results and Discussion

For each of the organophosphate pesticides in this investigation, the residues data obtained from the analytical method were mathematically transformed according to Equation 2 and, where possible, a first order decay model was fitted to the data using the statistical software R and following the equations above. Table 3 presents a summary of the WHP for the studied pesticides.

formulation dissolved in 50 L of water) compared to that used at the Al Soujah site (280 mL of 20% pesticide dissolved in 7 L of water). However, there is no information about the authenticity of the pesticide formulation. No statistical analysis could be carried out on the data relating to CPM.

3.2. Chlorpyrifos

The details of the regression model of chlorpyrifos presented in Table 4 indicates that there is an effect for the variables time and study area (Al-Hamah and Al Soujah), but there is no interaction between the variables.

Table 4. Details of the regression model of chlorpyrifos residue data (in natural logarithm scale) for dissipation time and study area.

Parameter	Statistical estimate	Standard Error	t value	Pr (> t)	Lower 95%	Upper 95%
Time 0 (intercept value)	4.73	0.19	24.76	<< 0.001	4.34	5.12
Dissipation time (slope value)	-0.209	0.016	-13.00	<< 0.001	-0.243	-0.1176
Study area (slope value)	-1.589	0.233	-6.83	<< 0.001	-2.069	-1.1096

That means it is possible to construct separate regression models for each of the study areas. However, it is acceptable to build a regression model with an average slope from both series [42] as shown in Figure 1A.




Figure 1. A) logarithmic decay of chlorpyrifos for the estimation of dissipation of chlorpyrifos; the fitted line was based on Al Soujah data; B) logarithmic decay of dimethoate for the estimation of dissipation of dimethoate and showing the effect of washing; C) logarithmic decay of diazinon for the estimation of dissipation of diazinon and showing linear and two compartment models; D) showing the modelled dissipation of the labile and persistent compartments for diazinon.

In this study the first sample of vine leaves was collected only after 60 minutes from pesticide application, due to manpower constraints and other operational factors. The volatile losses from leaves that may have occurred immediately after application can in fact be substantial [43]. Therefore, the initial application residue data, at time 0 hours, was estimated from the intercept of the regression model from the highest residue data set (Al Soujah). This conservative approach was taken to avoid overestimating the

half-lives. The intercept (i.e. time zero) was estimated from the constructed models. The initial pesticide residue loads for the two study locations were 114 mg/kg (Al Soujah series) (back transformed data for 4.73) and 23.22 mg/kg (back transformed data for 3.14) (Al Hamah series), see Table 4.

Although the initial loads seemed different between the study sites, from a statistical point of view there was no significant difference. The estimated half-life for chlorpyrifos was calculated using Equation 4, which gave an estimate of

3.3 days. Using the standard error of the slope (0.016), 95% confidence limits for the half-life were established at 2.9 – 3.9 days. The estimated half-life was similar for both data sets from each study area. These results for half-life values for chlorpyrifos are in line with previous ranges of 2.9-4 days as reviewed by Willies *et al.* [7] and Lu [10] in various foliage crops. The suggested withholding period (WHP) for chlorpyrifos was estimated using Equation 5. A key component of the estimate of the withholding period was the initial load. For chlorpyrifos this varied widely between the two study areas – each estimate having its own confidence limits. Several estimates of the initial load were, therefore, considered, including the average of the estimates, the maximum of the two estimates and the upper confidence limit of the upper estimate. The latter was chosen as a worst-case scenario, providing a safety margin when estimating the WHP. The initial load corresponded to 168.9 mg/kg. Depending on which MRL was used for the calculation, the WHP for chlorpyrifos, i.e. the time required to reduce the initial load of 168.9 mg/kg to 0.5 mg/kg if using the Codex MRL, or to 0.05 mg/kg if using the EU MRL for vine leaves as shown in Table 2, ranged from 26 to 37 days respectively. The WHP for chlorpyrifos is shown in Figure 1A as the time corresponding to the intersection point between the regression line and the MRL. This difference in WHP is significant and implicitly influences the agricultural practices in a relevant way. It also implies that vine leaves should not

be used for human or animal consumption for the entire duration of the WHP. Approximate confidence ranges for the withholding period (based on the initial load of 168.9 mg/kg) were then 22.2-29.5 days (using Codex MRL) and 31.6-42.1 days (using EU MRL). These confidence limits can only be considered as theoretical extrapolations assuming that the dissipation behaviour of chlorpyrifos approximated a linear function, as samples of vine leaves were not collected at those times. Each collected sample of vine leaves was also washed using tap water and boiling water. In the case of chlorpyrifos, there was no statistically significant effect of washing the vine leaves with tap water or hot water. There was also no interaction between washing and the effect of dissipation in time. That means that washing the leaves did not decrease the chlorpyrifos load at any of the residue levels studied.

3.3. Dimethoate

In the case of dimethoate there was no statistically significant difference between the data sets from the two study areas but there was a large difference between the washing treatments. There was no interaction between the washing treatments and the holding time. Washing with water removed the same fraction of the pesticide over time. Details of the initial estimated loads are given in Table 5.

Table 5. Details of regression of dimethoate against washing type and time.

Washing	Estimate (ln)	Std.Error	Estimated initial residue level at t0 (mg/kg)	t value	Pr (>F)
No washing	3.34	0.18	28.2	18.13	<< 0.001
Tap water	2.79	0.18	16.30	15.15	<< 0.001
Hot water	0.85	0.18	2.3	4.62	<< 0.001
Slope	-0.242	0.013		-18.9	<< 0.001

There was no significant difference between the two study areas. One of the results from the hot water wash series in area 1 (Al Soujah) had an initial value comparable to that of the samples with no washing. This is most probably an outlier but was included in the analysis as there was no valid reason for rejecting it. An estimate of the half-life was obtained using Equation 4 which gave an estimate of 2.9 days. By using the standard error of the slope, 95% confidence limits for the half-life were 2.6 – 3.2 days. These

estimated values are in line with previous findings for which the half-lives for dimethoate were in the range of 0.8-7.2 days depending on the crop [43]. The time required to reduce the initial load of 28.2 mg/kg (back-transformed value of 3.34) to the MRL value was calculated according to Equation 5 and shown in Figure 1B. The WHP for unwashed leaves was estimated as 19 days to accomplish the Codex MRL and 33 days to reach the EU MRLs, with approximate confidence intervals for the WHP as shown in Table 6.

Table 6. Estimated confidence intervals (in days) for withholding period for dimethoate for washed and unwashed leaves. Lower confidence level (LCL) and upper confidence level (UCL).

Washing	With EU-MRL of 0.01 mg/kg			With Codex Alimentarius-MRL of 0.3 mg/kg		
	LCL	Estimate (days)	UCL	LCL	Estimate (days)	UCL
None	31	33	34	18	19	20
Tap water	29	31	32	16	17	17
Hot water	22	23	24	8	8	9

These differences in WHP demonstrate the need to adopt the agricultural practices that best protect consumers and the environment. The MRL is a conventional measure that needs to reflect the conditions of use under specific environmental conditions, hence the need to generate sufficient support data, from all regions in the world, that can help set and harmonize

MRLs in international meetings and bodies such as the Codex Alimentarius Commission. Cabras *et al.* [44] provided data on the dissipation of dimethoate on grapes, showing that after 8 days from treatment 80% of dimethoate was dissipated in the field. The final residue then remained constant in the following 3 weeks. As to the effect of

washing, details of the regression of dimethoate against the washing regimes and holding time (average slope) and the series is given in Table 6. Figure 1 B shows the effect of washing on the dissipation of dimethoate. There was a very significant effect of hot water washing with an effective reduction of 92% of the residue level by washing the leaves with hot water. There was also a reduction of the residue level by washing with tap water but that effect was not statistically significant. This result is in line with results presented in an extensive review by Kaushik et al. [28], which demonstrates that washing and cooking help eliminate most of the pesticides in different food commodities.

3.4. Diazinon

In the case of diazinon the linear regression model according to Equation 2 fitted the data very poorly (correlation $r^2 = 0.78$). Diazinon showed a rapid decomposition phase in the first day (s) followed by a slower decomposition phase (see Figure 1C). This observation and the difference in dissipation kinetics may be explained by the fact that pesticides applied to the vine leaves may be adsorbed, absorbed, altered, volatilized, or removed by water and the integrated results of these processes may be seen as an initial rapid decline in surface residues followed by a slower, asymptotic decrease [36]. Environmental factors, especially sunlight through photolysis, may cause a very

rapid dissipation of diazinon within days from pesticide application. In addition, some of the diazinon may be absorbed by the plant tissues, reaching dynamic equilibrium and degrading thereafter very slowly [45].

An alternative two compartment model was therefore used for diazinon, as discussed by Torabi et al. [45]. The model is described by Equation 7, and graphically shown in Figure 1D.

$$C_t = C_A \times \exp(-k_1 \times \text{time}) + C_B \times \exp(-k_2 \times \text{time}) \quad (7)$$

Where C_A was the initial amount of rapid phase (estimated at $t=0$) and k_1 the dissipation rate in the rapid dissipation phase and C_B the amount in the slow phase (estimated at time $t = \text{change of slope}$) and k_2 the dissipation rate in the slow dissipation (persistent) phase. The parameter estimates for Equation 7 are shown in Table 7.

Table 7. Details of linear regression of diazinon residue data (in natural logarithm scale) against dissipation time for fast and slow phases.

Phase	Intercept		Slope	
	Estimate	Std. Error	Estimate	Std. Error
Fast	2.109	0.311	-1.043	0.420
Slow	0.296	0.533	-0.135	0.039

The slope of the regression line is derived by taking the differential from Equation 7 and estimated as shown by Equation 8:

$$K_{\text{diss}} (\text{Slope}) = -C_A \times k_1 \times \exp(-k_1 \times t) - C_B \times k_2 \times \exp(-k_2 \times t) \quad (8)$$

The estimated half-life of diazinon varies with time since the initial application of pesticides as shown in Figure 2.




Figure 2. Changes in estimated half-life for diazinon with time since pesticide application.

These results are in line with previous findings by Torabi *et al.* [45], Willis *et al.* [43], where half-lives data for diazinon were in the range of 0.4-5.3 days depending on the crop. The estimation of the initial load of diazinon at time 0 (application time) was estimated in two different ways. A direct method was to sum the C_A and C_B terms from Equation 7 (Table 8) and this gave an estimate of 10.06 mg/kg.

Table 8. Details of two compartmental model for diazinon residue data. Indication of the regression estimates for the slow component.

Parameter	Estimate (mg/kg)	Std. Error
CA	8.34	2.99
CB	1.72	2.61
k1	1.53	1.01
k2	0.139	0.16

An alternative method was to use the linear approximation, using the sum of the back-transformed intercepts, as shown in Table 7, which gave an estimation of 9.58 mg/kg. The two values are not significantly different. The WHP was then estimated based on the two-compartment model using the slow component. This gave an estimate, using the Codex MRL, of 8.9 days with confidence intervals of 6.0-10.6 days. The estimated time to achieve the EU's MRL was 37.1 days. The latter value is outside the range of the sampling time and is therefore considered as an approximation only, and no confidence limits can be provided. It is important that estimates of WHP are based on residue data from a sampling regime that includes the estimated time as one cannot be sure that the behaviour of the pesticides in the field follows a linear trend. In relation to the washing of the leaves for diazinon, there was no statistically significant effect of washing the vine leaves with tap water or hot water. There was also no interaction between washing and the effect of time. Although the difference was not statistically significant, the no wash treatment had more pesticide, about 30% more than the cold wash. Kaushik *et al.* [28] reviewed food processing treatments such as washing and cooking, among others, and demonstrated that in most cases washing and cooking particularly lead to large reductions in pesticide residue levels. Angioni *et al.* [46] demonstrated that washing with tap water reduced the residues of azoxystrobin and fenhexamid but not pyrimethanil. Similarly, in relation to washing, the three pesticides for which data were collected in this study behaved in very different ways. For better comprehension of the observed behaviour, Table 2 presents selected properties of the three studied pesticides. Chlorpyrifos is not a systemic insecticide [1], however it has a high Kow/Koc and would have been strongly bound to the cuticles of the leaves, so there would have been very little wash-off. This is in line with studies by Teixeira *et al.* [47] and Buschhaus *et al.* [48] on the composition of the epicuticular composition of leaves and their water barrier action as well as the mobility of pesticides in the plant where lipophilicity and concentration of the active ingredient are the driving forces

in the transfer. Adjuvants in commercial formulations may also regulate the transfer processes. Ling *et al.* [49] came to similar conclusions for this pesticide. Dimethoate has a low Kow/Koc and hence the wash-off was effective – especially when hot water was used. The dissipation of dimethoate in time, and especially the persistent portion, may be explained by translocation of the residues into other plant compartments, since it is classified as a systemic pesticide [47]. This is also in line with studies by Cabras *et al.* [50], where he showed that dimethoate degraded rapidly during the first week after pesticide application but remained constant in the following two weeks. The intermediate Kow/Koc values for diazinon may have resulted in a fraction of the pesticide being gradually absorbed into a protected layer, i.e. the epicuticular wax of the leaves, which progressively increases in terms of yield during the surface development of the vine leaves as suggested by Buschhaus *et al.* [48] and previously by Baker *et al.* [51], Torabi *et al.* [45], Willis *et al.* [7], Lu *et al.* [10]. The results from the present study agree with previous data for OPs. Willis *et al.* [7] reported average values of half-life for OPs of 3.0 ± 2.7 days for foliage. Cabras *et al.* [50] reported half-lives for OPs in grapes, wine and their processing products ranging between 0.97 and 3.84 days. Angioni *et al.* [22] showed decline curves for boscalid that do not follow a linear decay curve and stressed the idea that field agricultural practices critically influence the decrease or the disappearance of pesticide residues. Fantke *et al.* [13] provided impressive information by reviewing 811 publications in the scientific literature. Comparative assessment showed that that 95% of all half-lives, for all pesticide-plant species combinations, fall within the range of 0.6 and 29 days. Marin *et al.* [14] found that dissipation rates for selected pesticides (especially cyprodinil) were higher in the field at ambient temperature and with natural light than in cold conditions and in darkness. Chemical degradation caused by high temperatures and solar radiation is an important factor in pesticide dissipation kinetics. In the study presented here, environmental conditions possibly favoured chemical degradation in the field, which would have contributed to the short half-life of the studied OP pesticides.

4. Conclusions

Vine leave samples were collected over a 21-day period and analysed to assess the decay of four OPs, chlorpyrifos-methyl, chlorpyrifos, diazinon and dimethoate. Analysis of the data provided an indication of the dissipation rates, the half-lives and the WHPs. The three pesticides that could be evaluated behaved in very different ways. For diazinon, regression analysis could not fit the dissipation data to a first order decay model. The estimate of the half-life and withholding time for diazinon depended on the model used. The results from the present study are in line with previous data for organophosphorus pesticides. As part of the study

vine leaves were also washed with cold and boiling tap water. The data indicate that washing with hot water removed approximately 92% of the dimethoate residues. This offers a practical method of decreasing the intake of that pesticide and directly protects consumers of, for example, stuffed leaves. To conclude, the reported study can be replicated and applied in field monitoring, with the objective of gaining information on the results of current agricultural practices, thus contributing to set MRLs under the conditions of use in countries worldwide.

Acknowledgements

This work was supported by IAEA technical cooperation project SYR/5/023.

Appendix

The study areas are presented in Figures A1 and A2. They are Al Soujah (Yafour) area, located 23 km west of Damascus, and the Al-Hamah area located 14 km north west of Damascus respectively.



Figure A1. Al Soujah area. Vineyard surrounded by olive trees. Located west of Yafour.



Figure A2. Al Hamah area, a river crosses the area.

Table A1 gives the list of the pesticides, their retention times, the transitions used for quantitation and confirmation together with the optimised collision energies (CE) and the corresponding retention time windows in the multiple reaction monitoring (MRM) acquisition method obtained using a gas chromatograph coupled to tandem mass spectrometry (GC-MS/MS).

Table A1. List of pesticides, their retention times, the transitions used for quantitation and confirmation together with the optimised collision energies (CE) and the corresponding retention time windows in the multiple reaction monitoring (MRM) acquisition method.

Pesticides	MRM Time segment	RT (min)	Quantifier transition 1	CE 1 (V)	Qualifier transition 2	CE2 (V)
Chlorpyrifos	11	12.0	196.9 → 169.0	15	198.9 → 171.0	15
Chlorpyrifos-methyl	8	10.8	285.9 → 92.9	25	287.9 → 92.9	25
Diazinon	6	9.5	137.1 → 84.0	10	137.1 → 54.0	20
Dimethoate	4	8.6	86.9 → 46.0	15	92.9 → 63.0	10
Internal standards						
Sulfotep	3	8.0	201.8 → 145.9	10	201.8 → 81.9	25
Triphenyl phosphate	19	15.3	326.0 → 325.0	5	214.9 → 168.1	15

Method validation information for each pesticide including method limit of quantitation (LOQ), and the precision data obtained during method validation are presented in Table A2. The limit of quantitation (LOQ) of the method was the lowest fortification level of the validation meeting the method performance acceptability criteria and it was obtained from recovery studies.

Table A2. Method validation information for each pesticide including method limit of quantitation (LOQ), average recovery and intermediate precision data (repeatability and reproducibility relative standard deviation) at 0.01 mg/kg level.

Pesticide	LOQ (mg/kg)	Average recovery (%)	Repeatability RSD (%) at 0.01 mg/kg	Reproducibility RSD (%) at 0.01 mg/kg
Chlorpyrifos	0.01	83	15	16
Chlorpyrifos-methyl	0.01	81	13	15
Diazinon	0.01	82	13	14
Dimethoate	0.01	81	9	9

References

- E.; Gras, N.; Navarrete, D.; Lillo, P., Eds.; Pontificia Universidad Católica de Chile: Santiago de Chile, Chile, 2015; 72 pp.
- [1] Food and Agriculture Organization. FAO specifications and evaluations for agricultural pesticides 2018. www.fao.org/agriculture/crops/thematic-sitemap/theme/pests/jmps/ps-new/en/#D (accessed Jan 2018).
- [2] Maia, M.; Monteiro, F.; Sebastiana, M.; Marques, A. P.; Ferreira, A. E. N.; Freire, A. P.; Cordeiro, C.; Figueiredo, A.; Silva, M. S. Metabolite extraction for high-throughput FTICR-MS-based metabolomics of grapevine leaves. *EuPA Open Proteomics* 2016, 12, 4-9.
- [3] Tsiropoulos, N. G.; Miliadis, G. E.; Likas, D. T.; Liapis K. Residues of spiroxamine in grapes following field application and their fate from vine to wine. *J Agric Food Chem.* 2005, 53 (26), 10091-6.
- [4] Grimalt, S.; Dehouck, P. Review of analytical methods for the determination of pesticide residues in grapes. *J Chromatogr A* 2016, 12, 1433, 1-23.
- [5] Del Carlo, M.; Mascini, M.; Pepe, A.; Compagnone, D.; Mascini, M. Electrochemical Bioassay for the Investigation of Chlorpyrifos-methyl in Vine Samples. *J. Agric. Food Chem.* 2002, 50 (25), 7206–7210.
- [6] Arora, P. K.; Jyot, G.; Singh, B.; Singh Battu, R.; Singh, B.; Singh Aulakh, P. Persistence of Imidacloprid on Grape Leaves, Grape Berries and Soil. *Bull. Environ. Contam. Toxicol.* 2009, 82, 239–242.
- [7] Willis, G. H.; Mc Dowell, L. L. Pesticide persistence on foliage. *Reviews of environmental contamination and toxicology* 1987, 100, 23-73.
- [8] Pihlström, T. Analysis of pesticide residues in food using ethyl acetate extraction and detection with GC-MS/MS and LC-MS/MS (SweET). In *Proceedings of the 5th Latin American Pesticide Residues Workshop, food and environment, Cortes*; E.; Gras, N.; Navarrete, D.; Lillo, P., Eds.; Pontificia Universidad Católica de Chile: Santiago de Chile, Chile, 2015; 72 pp.
- [9] Montasser, M. R.; Mahmoud, H. A. Alexandria Science Exchange Journal 2009. www.alexexch.org/File/2009003001/En/2087.pdf (accessed Jun 2018).
- [10] Lu, M.-X., Jiang, W. W., Wang, J.-L., Liu, X.-J., Yu, X.-Y. Persistence and dissipation of chlorpyrifos in brassica chinensis, lettuce, celery, asparagus lettuce, eggplant, and pepper in a greenhouse. *PLoS ONE*, 2014, doi: 10.1371/journal.pone.0100556.
- [11] Food and Agriculture Organization and World Health Organization. Pesticide residues in food, 2016. www.fao.org/3/a-i5693e.pdf (accessed Jun 2018).
- [12] World Health Organization. JMPR toxicological monographs, 2017. www.who.int/foodsafety/publications/jmpr-monographs/en/ (accessed Jun 2018).
- [13] Fantke, P.; Juraske, R. Variability of pesticides dissipation half-lives in plants. *Environ. Sci. Technol.* 2013, 47, 3548-3562.
- [14] Marin, A.; Oliva, J.; Garcia, C.; Navarro, S.; Barba, A. Dissipation rates of cyprodinil and fludioxonil in lettuce and table grape in the field and under cold storage conditions. *J Agric Food Chem.* 2003, 51 (16), 4708-11.
- [15] Abdelraheem, E., Arief, M., Mohammad, S., G., Jiang, W. A Safety assessment of chromafenozide residue level with decline study on tomato in Egypt. *Environ Monit Assess.*, 2017, 189, 4, pp.180. doi: 10.1007/s10661-017-5894-6.
- [16] Hongfang, L., Xinze, L., Yecheng, M., Kyongjin, P., Jiye, H.. Residue analysis and dietary exposure risk assessment of tebufenozide in stem lettuce (*Lactuca sativa* L. var. angustana Irish). *Food and Chemical Toxicology*, 2018, 120, pp. 64-70. doi.org/10.1016/j.fct.2018.06.057.

- [17] Jie, L., Muhammad, R., Jiangwei, Q., Meiyang, H., Guohua, Z. Dissipation and metabolism of tebufenozide in cabbage and soil under open field conditions in South China, *Ecotoxicology and Environmental Safety*, 2016, 134, pp.204-212, doi.org/10.1016/j.ecoenv.2016.09.002.
- [18] Xiaoxin, C., Sheng, H., Yimen, G., Yecheng, M., Jiye, H., Xiao, L. Dissipation behavior, residue distribution and dietary risk assessment of field-incurred boscalid and pyraclostrobin in grape and grape field soil via MWCNTs-based QuEChERS using an RRLC-QqQ-MS/MS technique. *Food Chemistry*, 2018. doi.org/10.1016/j.foodchem.2018.08.136.
- [19] Soliman, A., S., Helmy, R., M., A., Nasr, I., N., Mahmoud, H., A., Jiang, W. Behavior of Thiophanate Methyl and Propiconazole in Grape and Mango Fruits Under the Egyptian Field Conditions. *Bull Environ Contam Toxicol.*, 2017, 98, 5, pp.720-725. doi: 10.1007/s00128-017-2066-x.
- [20] Liu J, Rashid M, Qi J, Hu M, Zhong G. Dissipation and metabolism of tebufenozide in cabbage and soil under open field conditions in South China. *Ecotoxicol Environ Saf.*, 2016, 134, pp. 204-212. doi:10.1016/j.ecoenv.2016.09.002.
- [21] Besil, N.; Pérez-Parada, A.; Cesio, V.; Varela, P.; Rivas, F., Heinzen H. Degradation of imazalil, orthophenylphenol and pyrimethanil in Clementine mandarins under conventional postharvest industrial conditions at 4°C. *Food Chem.* 2016, 1 (194), 1132-7.
- [22] Angioni; A.; Dedola, F.; Garau, V. L.; Schirra, M.; Caboni, P. Fate of iprovalicarb, indoxacarb, and boscalid residues in grapes and wine by GC-ITMS analysis. *J Agric Food Chem.* 2011, 59 (12), 6806-12.
- [23] Cabras, P.; Garau, V., L.; Filippo, M.; Pirisi, F. M.; Cubeddu, M.; Cabitza, F.; Spanedda, L. Fate of some insecticides from vine to wine. *J. Agric. Food Chem.* 1995, 43 (10), 2613–2615.
- [24] Navarro, S.; Oliva, J.; Navarro, G.; and Barba, A. Dissipation of chlorpyrifos, fenarimol, mancozeb, metalaxyl, penconazole, and vinclozolin in grapes. *American Journal of Enology and Viticulture* 2001, 52, 35–40.
- [25] Cus, F.; Basa Cesnik, H.; Bolta, S. V.; Gregorcic, A. Pesticide residues in grapes and during vinification process, *Food Control* 2010, 21 (11), 1512-1518.
- [26] Australian Government. *Agricultural and Veterinary Chemicals Code Act 1994*, latest edition 2016. www.legislation.gov.au/Series/C2004A04723 (accessed Jun 2018).
- [27] Amvrazi, E. G. Fate of Pesticide Residues on Raw Agricultural Crops after Postharvest Storage and Food Processing to Edible Portions. 2011. www.cdn.intechopen.com/pdfs/13027/InTech-Fate_of_pesticide_residues_on_raw_agricultural_crops_after_postharvest_storage_and_food_processing_to_edible_portions.pdf (accessed June 2018).
- [28] Kaushik, G.; Satya, S.; Naik, S. N. Food processing a tool to pesticide residue dissipation – A review. *Food Res. Int.* 2009, 42, 26-40.
- [29] Holland P. T., Hamilton D., Ohlin B., Skidmore M. W. Effects of storage and processing on pesticide residues in plant products. *Pure Appl. Chem.* 1994, 66, 335-356.
- [30] Maestroni, B.; Alnaser, A. Abu; Ghanem, I.; Islam, M.; Cesio, V.; Heinzen, H.; Kelly, S.; and Cannavan, A. Validation of an analytical method for the determination of selected pesticide residues in vine leaves by GC-MS/MS. *J. Agric. Food Chem.* 2018, DOI: 10.1021/acs.jafc.8b00453.
- [31] Lewis, K.; and Tzilivakis, J. Development of a data set of pesticide dissipation rates in/on various plant matrices for the Pesticide Properties Data Base (PPDB). *Data* 2017, 2 (3), 28.
- [32] Galiotta, G.; Egana, E.; Gemelli, F.; Maeso, D.; Casco, N.; Conde, P.; Nunez, S. Pesticide dissipation curves in peach, pear and tomato crops in Uruguay. *J Environ Sci Health B.* 2011, 46 (1), 35-40.
- [33] Yajie, C., Mingcheng, G., Xingang, L., Jun, X., Fengshou, D., Xiaohu, W., Baotong, L., Yongquan, Z. Determination and dissipation of afidopyropen and its metabolite in wheat and soil using QuEChERS-UHPLC-MS/MS. *J. Sep. Science*, 2018, 41, 7, pp. 1674-1681, doi.org/10.1002/jssc.201700773.
- [34] Xiaokang, A., Jun, X., Fengshou, D., Xingang, L., Xiaohu, W., Ran, W., Yongquan, Z. Simultaneous determination of broflanilide and its metabolites in five typical Chinese soils by a modified quick, easy, cheap, effective, rugged, and safe method with ultrahigh - performance liquid chromatography and tandem mass spectrometry, *J. Sep. Science*, 2018, doi.org/10.1002/jssc.201800631.
- [35] Huan, Z., Luo, J., Xu, Z., Xie, D. Residues, dissipation, and risk assessment of spinosad in cowpea under open field conditions. *Environ. Monit. Assess.* 2015, 187, 11, pp. 706. doi: 10.1007/s10661-015-4942-3.
- [36] Saber, A., N, Malhat, F., M, Badawy, H., M, Barakat, D., A. Dissipation dynamic, residue distribution and processing factor of hexythiazox in strawberry fruits under open field condition. *Food Chem.* 2016, 196, pp. 1108-1116. doi:10.1016/j.foodchem.2015.10.052.
- [37] Malhat, F., Badawy, H., M., A., Barakat, D., A., Saber, A., N. Residues, dissipation and safety evaluation of chromafenozide in strawberry under open field conditions. *Food Chemistry*, 2014, 152, pp. 18-22. doi: 10.1016/j.foodchem.2013.11.110.
- [38] Malhat, F., El-Mesallamy, A., Assy, M., Madian, W., Loutfy, N., M., Tawfic, M., A. Residues, half-life times, dissipation, and safety evaluation of the acaricide fenpyroximate applied on grapes, *Toxicological & Environmental Chemistry*, 2013, 95, 8, 1309-1317, DOI: 10.1080/02772248.2013.877245.
- [39] R Core Team, 2017. <https://www.r-project.org/> (accessed Jun 2018).
- [40] Paramasivam, M.; Deepa, M; Selvi, C.; and Chandrasekaran, S. Dissipation kinetics and safety evaluation of tebuconazole and trifloxystrobin in tea under tropical field conditions. *Food Additives & Contaminants: Part A* 2017, 34 (12), 2155-2163.
- [41] Upton, G., Cook, I. *Oxford Dictionary of Statistics* (3rd Edition) Oxford Univ. Press: Oxford, UK, 2014.
- [42] Efron, B.; Tibshirani, R. *An Introduction to the Bootstrap*. Chapman and Hall / CRC: New York, 1994.
- [43] Willis, G. H.; Mc Dowell, L. L.; Southwick, L. M.; Smith, S. Toxaphene, methyl parathion, and fenvalerate disappearance from cotton foliage in the Mid South. *J. Environ. Qual.* 1985, 14, 446-450.
- [44] Cabras, P.; Garau, V. L.; Melis, M.; Pirisi, F. M.; Cubeddu, M.; Cabitza F. Residui di dimetoato e clorpirifos nell'uva e nel vino. *Atti Giornate Fitopatologiche* 1994, 1, 27-32.

- [45] Torabi, E.; Talebi, K. Diazinon residues and degradation kinetics for grapes under field conditions. *J Environ Sci Health B*. 2013, *48* (4), 260-5.
- [46] Angioni, A.; Schirra, M.; Garau, V. L.; Melis, M.; Tuberoso, C. I. G.; Cabras, P. Residues of azoxystrobin, fenhexamid and pyrimethanil in strawberry following field treatments and the effect of domestic washing. *Food Addit. Contam.* 2004, *21*, 1065-1070.
- [47] Teixeira, M.; Aguiar, A.; Afonso, C.; Alves, A.; Bastos, M. Comparison of pesticides levels in grape skin and in the whole grape by a new liquid chromatographic multiresidue methodology. *Analytica Chimica Acta*. 2004, *513*, 333-340.
- [48] Buschhaus, C.; and Jetter, R. Composition and Physiological Function of the Wax Layers Coating Arabidopsis Leaves: β -Amyrin Negatively Affects the Intracuticular Water Barrier. *Plant Physiology* 2012, *160* (2), 1120-1129.
- [49] Ling, Y.; Wang, H.; Yong, W.; Zhang, F.; Sun, L.; Yang, M. L.; Wu, Y. N.; Chu, X. G. The effects of washing and cooking on chlorpyrifos and its toxic metabolites in vegetables. *Food Control* 2011, *22*, 54-58.
- [50] Cabras, P.; and Angioni, A. Pesticide Residues in Grapes, Wine, and Their Processing Products. *J. Agric. Food Chem.* 2000, *48* (4), 967-973.
- [51] Baker, E. A.; Hunt, G. M. Developmental changes in leaf epicuticular waxes in relation to foliar penetration. *New Phytologist* 1981, *88*, 731-747.